

Constraints on Higher-Order Perturbative Corrections in $b \rightarrow u$ Semileptonic Decays from Residual Renormalization-Scale Dependence

M.R. Ahmady¹, F.A. Chishtie², V. Elias², A.H. Fariborz³,
D.G.C. McKeon², T.N. Sherry⁴, T.G. Steele⁵

February 1, 2008

Abstract

The constraint of a progressive decrease in residual renormalization scale dependence with increasing loop order is developed as a method for obtaining bounds on unknown higher-order perturbative corrections to renormalization-group invariant quantities. This technique is applied to the inclusive semileptonic process $b \rightarrow u\bar{\nu}_\ell\ell^-$ (explicitly known to two-loop order) in order to obtain bounds on the three- and four-loop perturbative contributions that are not accessible via the renormalization group. Combining this technique with the principle of minimal sensitivity, we obtain an estimate for the perturbative contributions to $\Gamma(b \rightarrow u\bar{\nu}_\ell\ell^-)$ that incorporates theoretical uncertainty from as-yet-undetermined higher order QCD corrections.

A variety of techniques have been developed for obtaining estimates of higher-loop corrections in perturbative QCD, including the principle of minimal sensitivity (PMS) [1], the fastest apparent convergence (or effective charges) approach [2], nonabelianization methods [3], Padé approximations [4], and renormalization-group supplemented Padé approximants [5]. Of particular relevance is the PMS ansatz which proposes that renormalization-scheme independence, which at lower-loop levels translates into renormalization-scale (μ) independence, provides the optimum perturbative prediction of a QCD observable.

In a particular renormalization scheme (such as $\overline{\text{MS}}$), the minimal sensitivity principle identifies the appropriate choice of renormalization scale μ for a physical observable as the value at which the observable is independent of μ , providing a method for dealing with the residual renormalization scale dependence that exists in a perturbative calculation truncated to any given order. In explicit calculations, such residual scale dependence decreases as higher-order corrections are included; if a renormalization-group-invariant quantity Γ is truncated at order $(\alpha/\pi)^n$, one can easily demonstrate that $d\Gamma/d\mu$ is order $(\alpha/\pi)^{n+1}$. Consequently, perturbative predictions become virtually μ -independent at sufficiently high orders. For example, in inclusive semileptonic $b \rightarrow u$ decays, a clear progressive flattening of the decay rate as a function of μ is observed in the explicit calculations up to two-loop order [6], and this property persists when a Padé estimate of the three-loop correction is included [7].

In this paper we demonstrate for semileptonic $b \rightarrow u$ decays that the progressive decrease in renormalization-scale dependence with increasing loop-order places bounds on unknown higher-order perturbative coefficients. Combined with minimal-sensitivity, the prediction of the decay rate devolving from these bounds leads to an estimate of higher-order perturbative corrections to the decay rate.

The perturbative contributions to the inclusive semileptonic $b \rightarrow u$ decay rate $\Gamma(b \rightarrow u\bar{\nu}_\ell\ell^-)$ may be expressed

¹Department of Physics, Mount Allison University, Sackville, NB E4L 1E6, Canada

²Department of Applied Mathematics, The University of Western Ontario, London, ON N6A 5B7, Canada

³Dept. of Mathematics/Science, State Univ. of New York Institute of Technology, Utica, NY 13504-3050, USA

⁴Department of Mathematical Physics, National University of Ireland, Galway, Ireland

⁵Department of Physics & Engineering Physics, University of Saskatchewan, Saskatoon, SK S7N 5E2, Canada

as [6]

$$\begin{aligned} \frac{1}{K} \Gamma(\mu, m_b(\mu), x(\mu)) = m_b^5(\mu) & \left(1 + [a_0 - a_1 \log(w)] x(\mu) + [b_0 - b_1 \log(w) + b_2 \log^2(w)] x^2(\mu) \right. \\ & + [c_0 - c_1 \log(w) + c_2 \log^2(w) - c_3 \log^3(w)] x^3(\mu) \\ & \left. + [d_0 - d_1 \log(w) + d_2 \log^2(w) - d_3 \log^3(w) + d_4 \log^4(w)] x^4(\mu) + \dots \right) \end{aligned} \quad (1)$$

where

$$x(\mu) \equiv \frac{\alpha_s(\mu)}{\pi} \quad , \quad w = w(\mu, m_b(\mu)) \equiv \frac{m_b^2(\mu)}{\mu^2} \quad , \quad K \equiv \frac{G_F^2 |V_{ub}|^2}{192\pi^3} \quad . \quad (2)$$

In the $\overline{\text{MS}}$ scheme in which m_b is identified with a scale-dependent (running) b -quark mass for four or five active flavours, the one- and two-loop order coefficients within (1) are given by [6]

$$\begin{aligned} \text{all } n_f : \quad a_0 &= 4.25360 \quad , \quad a_1 = 5 \quad , \\ n_f = 5 : \quad b_0 &= 26.7848 \quad , \quad b_1 = 36.9902 \quad , \quad b_2 = 17.2917 \quad , \\ n_f = 4 : \quad b_0 &= 25.7547 \quad , \quad b_1 = 38.3935 \quad , \quad b_2 = 17.7083 \quad . \end{aligned} \quad (3)$$

The running mass $m_b(\mu)$ appearing in (1) is also known to full four-loop order [9]:

$$m_b[x(\mu)] = m_b[x(\mu_0)] \frac{c[x(\mu)]}{c[x(\mu_0)]} \quad , \quad (4)$$

$$n_f = 4 : \quad c[x] = x^{12/25} [1 + 1.01413x + 1.38920x^2 + 1.09052x^3 + \dots] \quad , \quad (5)$$

$$n_f = 5 : \quad c[x] = x^{12/23} [1 + 1.17549x + 1.50071x^2 + 0.172486x^3 + \dots] \quad . \quad (6)$$

Renormalization-group invariance of the decay rate Γ determines a subset of the unknown three-loop c_k coefficients and four-loop d_k coefficients. From the renormalization-group equation

$$\begin{aligned} 0 &= \mu^2 \frac{d}{d\mu^2} \Gamma(\mu, m_b(\mu), x(\mu)) \\ &= \left[\mu^2 \frac{\partial}{\partial \mu^2} - (\gamma_0 x + \gamma_1 x^2 + \gamma_2 x^3 + \dots) m_b \frac{\partial}{\partial m_b} - (\beta_0 x^2 + \beta_1 x^3 + \dots) \frac{\partial}{\partial x} \right] \Gamma \quad , \end{aligned} \quad (7)$$

we find the following expressions for terms proportional to x^3 and x^4 in the final line of (7):

$$c_1 = 2b_0\beta_0 + a_0\beta_1 + \gamma_0(5b_0 - 2b_1) + \gamma_1(5a_0 - 2a_1) + 5\gamma_2 \quad , \quad (8)$$

$$c_2 = \frac{1}{2} [2b_1\beta_0 + a_1(\beta_1 + 5\gamma_1) + \gamma_0(5b_1 - 4b_2)] \quad , \quad (9)$$

$$c_3 = \frac{b_2}{3} (2\beta_0 + 5\gamma_0) \quad , \quad (10)$$

$$d_1 = \beta_2 a_0 + 2\beta_1 b_0 + 3\beta_0 c_0 + \gamma_0(5c_0 - 2c_1) + \gamma_1(5b_0 - 2b_1) + \gamma_2(5a_0 - 2a_1) + 5\gamma_3 \quad (11)$$

$$d_2 = \frac{1}{2} \beta_2 a_1 + \beta_1 b_1 + \frac{3}{2} \beta_0 c_1 + \gamma_0 \left(\frac{5}{2} c_1 - 2c_2 \right) + \gamma_1 \left(\frac{5}{2} b_1 - 2b_2 \right) + \frac{5}{2} \gamma_2 a_1 \quad (12)$$

$$d_3 = \beta_0 c_2 + \frac{2}{3} \beta_1 b_2 + \gamma_0 \left(\frac{5}{3} c_2 - 2c_3 \right) + \frac{5}{3} \gamma_1 b_2 \quad (13)$$

$$d_4 = \frac{3}{4} \beta_0 c_3 + \frac{5}{4} \gamma_0 c_3 \quad (14)$$

Upon substitution of (3) and the four-loop $\overline{\text{MS}}$ results for the β function [8] and anomalous mass dimension γ [9] into (8–14), we obtain the following numerical values for these higher-loop coefficients:

$$c_1^{(4)} = 263.839, \quad c_2^{(4)} = 194.234, \quad c_3^{(4)} = 54.1087 \quad (15)$$

$$d_1^{(4)} = 11.25c_0^{(4)} + 103.081, \quad d_2^{(4)} = 1580.26, \quad d_3^{(4)} = 765.844, \quad d_4^{(4)} = 152.181 \quad (16)$$

$$c_1^{(5)} = 249.592, \quad c_2^{(5)} = 178.755, \quad c_3^{(5)} = 50.9145 \quad (17)$$

$$d_1^{(5)} = 10.75c_0^{(5)} - 8.28683, \quad d_2^{(5)} = 1376.68, \quad d_3^{(5)} = 667.838, \quad d_4^{(5)} = 136.833 \quad (18)$$

where the superscript denotes the number of active flavours (either $n_f = 4$ or $n_f = 5$). In the energy region spanning the threshold between $n_f = 4$ and $n_f = 5$ there are four higher-loop parameters that remain undetermined: $\{c_0^{(4)}, c_0^{(5)}, d_0^{(4)}, d_0^{(5)}\}$. However, continuity of the decay rate at this threshold in conjunction with the coupling-constant and quark-mass threshold matching conditions imposed at $\mu = m_b$ [10],¹

$$x^{(4)}(m_b) = x^{(5)}(m_b) \left[1 + 0.1528 \left[x^{(5)}(m_b) \right]^2 + 0.633 \left[x^{(5)}(m_b) \right]^3 \right] \quad (19)$$

$$m_b^{(4)}(m_b) = m_b^{(5)}(m_b) \left[1 + 0.2060 \left[x^{(5)}(m_b) \right]^2 + 1.9464 \left[x^{(5)}(m_b) \right]^3 \right], \quad (20)$$

effectively reduces this set to two unknown parameters. Using the benchmark value $\alpha_s(M_z) = 0.119$ [11], $m_b = 4.2 \text{ GeV}$ [12], and the four-loop β function, we first find that

$$x^{(5)}(4.2 \text{ GeV}) = 0.07261, \quad x^{(4)}(4.2 \text{ GeV}) = 0.07269 \quad (21)$$

$$m_b^{(4)}(4.2 \text{ GeV}) = 4.208 \text{ GeV}. \quad (22)$$

Imposing continuity of the decay rates to $\mathcal{O}(x^3)$ at this flavour threshold then yields

$$c_0^{(5)} = 1.012c_0^{(4)} + 15.66. \quad (23)$$

Such a continuity condition cannot be extended to $\mathcal{O}(x^4)$, because the first unknown term in the mass threshold condition (20) contributes at $\mathcal{O}(x^4)$ to the d_0 coefficient. In the absence of this information, we assume

$$d_0^{(4)} \approx d_0^{(5)}, \quad (24)$$

reflecting the near-equivalence of the known coefficients $\{d_2, d_3, d_4\}$ for four and five active flavours.² Thus the RG analysis combined with continuity of the decay rate at flavour thresholds implies that the four-loop expression (1) effectively has only two undetermined parameters $\{c_0^{(4)}, d_0^{(4)}\}$ in the energy range spanning the threshold between $n_f = 4$ and $n_f = 5$.

The decay rate Γ is a truncated perturbation series which necessarily exhibits residual renormalization scale dependence. In general if Γ is known to $\mathcal{O}(x^n)$, then $d\Gamma/d\mu = \mathcal{O}(x^{n+1})$. Consequently, for any reasonable perturbation series, the residual scale dependence diminishes with increasing loop order. A measure of the residual scale dependence in the natural energy region³ $m_b/2 < \mu < 2m_b$ is provided by the deviation of Γ from its average value

$$\chi^2 = \frac{1}{\frac{3}{2}m_b} \int_{m_b/2}^{2m_b} \left(\frac{\Gamma(\mu)}{\langle \Gamma \rangle} - 1 \right)^2 d\mu, \quad (25)$$

¹In the $\overline{\text{MS}}$ scheme the scale $\mu = m_b$ is defined by $m_b(m_b) = m_b$.

²In (20), if the coefficient of $[x^{(5)}(m_b)]^4$ were 20, reflecting a factor of ten increase between the two previous orders, the difference between $d_0^{(4)}$ and $d_0^{(5)}$ would be approximately 150, which is small compared to the scales of d_0 we obtain in our analysis below.

³This energy region is exactly that considered in [6] where the residual scale dependence of the two-loop decay rate was originally considered.

where the pre-factor of $3m_b/2$ leads to a dimensionless χ^2 , and where $\langle\Gamma\rangle$ is the average value of Γ over the same energy interval as (25):

$$\langle\Gamma\rangle = \frac{1}{\frac{3}{2}m_b} \int_{m_b/2}^{2m_b} \Gamma(\mu) d\mu \quad . \quad (26)$$

The progressive decrease in Γ 's scale dependence implies that χ^2 must decrease as the loop order of Γ is increased. However, since at three-loop order Γ (and hence χ^2) will depend on the parameter $c_0^{(4)}$, and at four-loop order will depend on $\{c_0^{(4)}, d_0^{(4)}\}$, the progressive decrease in χ^2 as loop-order increases necessarily provides constraints on these unknown higher-loop parameters.

To obtain such constraints, we use the central values $\alpha_s(M_z) = 0.119$ [11], $m_b = 4.2$ GeV [12], the four-loop β and γ functions, and the threshold matching conditions (19), (20) to evaluate $\alpha(\mu)$ and $m_b(\mu)$. We then utilize discretization to evaluate the integrals (25) and (26) and obtain χ^2 values shown in Figure 1. The two-loop result is by definition independent of $c_0^{(4)}$; the three-loop result depends on $c_0^{(4)}$ as displayed in Figure 1. The requirement that the three-loop χ^2 is less than the two-loop χ^2 is satisfied provided

$$-150 < c_0^{(4)} < 290 \quad . \quad (27)$$

The Padé estimate $c_0^{(4)} = 188$ of [7] is well within this interval. Combining the result (27) with (23) we find the following constraint on $c_0^{(5)}$:

$$-136 < c_0^{(5)} < 309 \quad , \quad (28)$$

which, in the notation $c_0^{(5)} = 200 + \Delta$, corresponds to $-336 < \Delta < 109$ that includes the range of estimates $-100 \leq \Delta \lesssim 80$ discussed in [13]. The optimal value of $c_0^{(4)}$, *i.e.*, the value that minimizes χ^2 , occurs at $c_0^{(4)} = 57$ ($c_0^{(5)} = 74$) corresponding most closely to the large- β_0 estimate ($\Delta = -100$) of [13, 14].

The progressive decrease in χ^2 to four-loop order restricts $\{c_0^{(4)}, d_0^{(4)}\}$ parameter space to the region indicated in Figure 2. Further insight into the allowed $d_0^{(4)}$ range can be obtained by employing the Padé estimate $c_0^{(4)} = 188$ [7] to determine a constant three-loop value for χ^2 . Upon comparison with the four-loop χ^2 dependence on the unknown parameter $d_0^{(4)}$, we find from Figure 3 that χ^2 continues to decrease with increasing order provided

$$-500 \lesssim d_0^{(4)} \lesssim 2500 \quad . \quad (29)$$

Although the range of parameter space in Figure 2 [and in (27)–(29)] might superficially be dismissed as too large to be of interest, it actually provides valuable control over higher-order corrections in the prediction of the actual phenomenological decay rate. If we extract the minimal-sensitivity prediction of the decay rate over the entire parameter space of Figure 2,⁴ we obtain the following range for the perturbative contributions to the inclusive semileptonic decay rate:

$$\frac{\Gamma}{K} = 2050 \pm 270 \text{ GeV}^5 \quad . \quad (30)$$

By contrast, the perturbative two-loop rate, obtained by truncation of the series (1) after its (fully known) $\mathcal{O}(x^2)$ contributions, does not exhibit minimal sensitivity at all in the $m_b/2 \leq \mu \leq 2m_b$ region, but is seen to decrease with increasing μ from 2053 GeV^5 to 1622 GeV^5 . Indeed, one unanticipated effect of including three- and four-loop coefficients, both known $\{c_{1-3}, d_{2-4}\}$ and unknown, is to ensure that minimal sensitivity occurs in the range $m_b/2 \leq \mu \leq 2m_b$ over almost all of the parameter space of Figure 2. This property is shown in Figure 4 which

⁴Minimal sensitivity occurs when $d\Gamma/d\mu = 0$ in the region $m_b/2 \leq \mu \leq 2m_b$. When there exists more than one critical point, we choose the one with the smallest second derivative, corresponding to the least sensitive choice.

compares the two- three- and four-loop contributions to the reduced rate for a typical point in the Fig. 2 parameter space.

The central value of the bound (30) corroborates the Padé estimate 2071 GeV^5 obtained in [7]. The $\pm 13\%$ uncertainty in the rate (30), obtained from the combined constraints of a progressive decrease in residual scale dependence and minimal sensitivity, is a genuine reflection of the uncertainty following from unknown higher-order QCD corrections. Such corrections, which typically are legislated away via series truncation after known-order contributions, cannot be disregarded as a source of additional uncertainty in predictions following from perturbative quantum field theory.

Acknowledgments: We are grateful for the hospitality of the KEK Theory Group, where this research was initiated with financial support from the International Opportunity Fund of the Natural Sciences and Engineering Research Council of Canada (NSERC) and the International Collaboration Programme 2001 of Enterprise Ireland.

References

- [1] P.M. Stevenson, Phys. Rev. D 23 (1981) 2916 and Phys. Lett. B 100 (1981) 61.
- [2] G. Grunberg, Phys. Lett. B 95 (1980) 70 and Phys. Rev. D 29 (1984) 2315.
- [3] D.J. Broadhurst, A.L. Kataev, C.J. Maxwell, Nucl. Phys. B 592 (2001) 247.
- [4] J. Ellis, M. Karliner, and M.A. Samuel, Phys. Lett. B 400 (1997) 176;
J. Ellis, I. Jack, D.R.T. Jones, M. Karliner, and M.A. Samuel, Phys. Rev. D 57 (1998) 2665.
- [5] V. Elias, T.G. Steele, F. Chishtie, R. Migneron, K. Sprague, Phys. Rev. D 58 (1998) 116007;
F. Chishtie, V. Elias, T.G. Steele, Phys. Rev. D 59 (1999) 105013;
M.R. Ahmady, F.A. Chishtie, V. Elias, A.H. Fariborz, D.G.C. McKeon, T.N. Sherry, T.G. Steele, hep-ph/0109061 (to appear, Phys. Rev. D).
- [6] T. van Ritbergen, Phys. Lett. B 454 (1999) 353.
- [7] M.R. Ahmady, F.A. Chishtie, V. Elias, and T.G. Steele, Phys. Lett. B 479 (2000) 201.
- [8] T. van Ritbergen, J.A.M. Vermaseren, and S.A. Larin, Phys. Lett. B 400 (1997) 379.
- [9] K.G. Chetyrkin, Phys. Lett. B 404 (1997) 161;
J.A.M. Vermaseren, S.A. Larin, T. van Ritbergen, Phys. Lett. B 405 (1997) 327.
- [10] K.G. Chetyrkin, B.A. Kniehl, M. Steinhauser, Nucl. Phys. B 510 (1998) 61.
- [11] Particle Data Group (D.E. Groom *et al*), Eur. Phys. J. C 15 (2000) 1.
- [12] A.H. Hoang, Phys. Rev. D 61 (2000) 034005 and D 59 (1999) 014039;
K.G. Chetyrkin, M. Steinhauser, Phys. Rev. Lett. 83 (1999) 4001.
- [13] K. Melnikov, T. van Ritbergen, Phys. Lett. B 482 (2000) 99.
- [14] M. Beneke and V.M. Braun, Phys. Lett. B 348 (1995) 513;
P. Ball, M. Beneke and V.M. Braun, Phys. Rev. D 52 (1995) 3929.




Figure 1: The quantity χ^2 as a function of $c_0^{(4)}$ (solid curve) for the three-loop contributions to Γ . The straight dashed line represents the two-loop χ^2 , and the criterion that the three-loop χ^2 is smaller than the two-loop χ^2 constrains $c_0^{(4)}$ to the region between the intersection points of the three- and two-loop curves.




Figure 2: The dots correspond to points within $\{c_0^{(4)}, d_0^{(4)}\}$ parameter space for which χ^2 decreases as loop-order increases from two to four. The sharp cutoff in the $c_0^{(4)}$ direction corresponds to the range obtained from Figure 1.




Figure 3: The quantity χ^2 as a function of $d_0^{(4)}$ (dashed curve) for the four-loop contributions to Γ after input of the Padé estimate $c_0^{(4)} = 188$. The straight solid line represents the three-loop χ^2 . The criterion that the four-loop χ^2 is smaller than three-loop χ^2 constrains $d_0^{(4)}$ to the region between the intersection points of the curves.




Figure 4: Two-loop (solid curve), three-loop (dotted curve) and four-loop (dashed-curve) values for the reduced rate Γ/K with $c_0^{(4)} = 80$ (corresponding to a central value $\Delta = 114$ from [13]) and $d_0^{(4)} = -500$ (central value in the Fig. 2 parameter space for the chosen $c_0^{(4)}$). Progressive decrease of residual scale dependence with increasing loop order is evident, as is the existence of PMS points for the three- and four-loop curves. Note the absence of a PMS point for the two-loop curve.